

# **Development of Risk-based Environmental Assessment and Remediation Guidelines for Salt Releases to Muskeg and Other Wetlands Settings**

Craig Harris, AECOM Canada  
GL908051

Moderately to highly saline groundwater co-occurs with most petroleum oil and natural gas deposits. Releases of saline “produced water” potentially affect peatland-forming wetlands (bogs and fens), particularly in temperate to sub-arctic circumpolar environments. This study quantifies vegetative and soil faunal responses to salinization of western Canadian boreal peatlands, towards the development of ecological risk-based remediation guidance.

Field data on the abundance (or percent cover) of vascular plants, bryophytes, and soil mesofauna were obtained in the summer of 2008 and 2009 from nine produced water release sites in Alberta and British Columbia. Research plots (1 m<sup>2</sup> quadrat and peat core sample sites) were established along salinity

gradients within fen (n=5), bog (n=3) and marsh (n=1) ecosystems arising from recent spills (mostly (predominantly fen sites), while a statistically significant relationship between species richness and the log10 of salt levels was different for a large marsh site in central Alberta. Marshes generally exhibit higher plant diversity than peatlands: Plots at the marsh site exhibited higher plant diversity at a given salinity than the fen sites, but the slope of the best-fit line describing the species richness – salt concentration relationship was similar to that of the fen sites.

The overall presence-absence data provide a clear indication of the salinity tolerance ranges of 31 dominant vascular plant and bryophytes species, including three commonly occurring species of sphagnum moss, rough bentgrass, black spruce, dwarf birch, Labrador tea, and dwarf bog cranberry. Some plants such as fireweed, Labrador tea, dwarf birch or aquatic sedge were observed to persist at soil salinities in excess of 6 – 8 mS/cm, while some species such as *Sphagnum girgenshoni* or common strawberry were not observed at soil salinities in excess of 3 to 4 mS/cm. The vast majority of observed plants and bryophytes (101 of 162 observed species) were observed in only one or a few of the 128 plots total. The salinity tolerances of these more rare species cannot be defined without use of an alternative methodology, since presence in any plot for rare species is ascribed firstly to 'chance' encounters and

secondarily to the salt levels in the co-located soils and interstitial water.

No appreciable relationship could be discerned between soil or interstitial salt levels and the abundance or composition of soil invertebrates (mesofauna) extracted from peat cores obtained from the same plots used to assess vegetative ecology. The absence of a discernible influence of plot salinity on mesofaunal abundance or composition is hypothesized to have resulted from microenvironmental variability in near-surface environments (both laterally and horizontally) that was not adequately captured by the sampling methods, and identification of mesofaunal taxa to only the family level or higher.

Occurrence data for each plant and bryophyte species that were observed in eight or more of the 127 plots assessed at field sites allow for estimation of frequency of occurrence statistics for each taxon along a ( $\log_{10}$ ) gradient of increasing salinity. The field data for Alberta peatland environments provide evidence of salinity ranges at which various plants may persist, at least for shorter time periods of approximately a half decade or less. Among the commonly occurring peatland sphagnidae, *S. magellanicum* was observed to be relatively salt tolerant, while *S. girgenshonii* and *S. fuscum* are relatively salt intolerant. Black spruce (*P. mariana*) is poorly tolerant to salinization; therefore, at historical release sites, it can be expected that the

majority of other peatland vegetation can potentially recolonize the disturbance site once *P. mariana* seedlings start to appear.

2007 UMA\_Risk Based Salinity Guidelines for Peatlands\_Literature Review

2008 UMA\_RiskBased Salt Releases to Muskeg Presentation

2007 UMA\_Risk Based Environmental Quality Guidelines for Salt Releases to Muskeg and Other Boreal Wetlands\_Proposal Report